

A THIRD-ORDER APÉRY-LIKE RECURSION FOR $\zeta(5)$

WADIM ZUDILIN (Moscow)

E-print math.NT/0206178

27 May 2002

Last revision: 06 July 2002

In 1978, R. Apéry [1], [2] has given sequences of rational approximations to $\zeta(2)$ and $\zeta(3)$ yielding the irrationality of each of these numbers. One of the key ingredient of Apéry's proof are second-order difference equations with polynomial coefficients satisfied by numerators and denominators of the above approximations. Recently, V. N. Sorokin [3] and this author [4], [5] have got independently a similar second-order difference equation for $\zeta(4)$. This recursion does not give diophantine approximations to $\zeta(4) = \pi^4/90$ proving its irrationality, however it presents an algorithm for fast calculation of this constant. The aim of this note is to highlight a possible generalization of the above results for the number $\zeta(5)$, the irrationality of which is not yet proved since nowadays.

1. Statement of the main result. Consider the difference equation

$$(n+1)^6 a_0(n) q_{n+1} + a_1(n) q_n - 4(2n-1) a_2(n) q_{n-1} - 4(n-1)^4 (2n-1)(2n-3) a_0(n+1) q_{n-2} = 0, \quad (1)$$

where

$$\begin{aligned} a_0(n) &= 41218n^3 - 48459n^2 + 20010n - 2871, \\ a_1(n) &= 2(48802112n^9 + 89030880n^8 + 36002654n^7 - 24317344n^6 - 19538418n^5 \\ &\quad + 1311365n^4 + 3790503n^3 + 460056n^2 - 271701n - 60291), \\ a_2(n) &= 3874492n^8 - 2617900n^7 - 3144314n^6 + 2947148n^5 + 647130n^4 \\ &\quad - 1182926n^3 + 115771n^2 + 170716n - 44541, \end{aligned}$$

and define its three linearly independent solutions $\{q_n\}$, $\{p_n\}$, and $\{\tilde{p}_n\}$ by the initial data

$$\begin{aligned} q_0 &= -1, \quad q_1 = 42, \quad q_2 = -17934, \quad p_0 = 0, \quad p_1 = \frac{87}{2}, \quad p_2 = -\frac{1190161}{64}, \\ \tilde{p}_0 &= 0, \quad \tilde{p}_1 = \frac{101}{2}, \quad \tilde{p}_2 = -\frac{344923}{16} \end{aligned}$$

(here and below a symbol $\{x_n\}$ means the sequence $\{x_n\}_{n=0}^\infty = \{x_0, x_1, x_2, \dots\}$).

Theorem 1. *The sequences*

$$\ell_n = q_n \zeta(5) - p_n, \quad \tilde{\ell}_n = q_n \zeta(3) - \tilde{p}_n, \quad n = 0, 1, 2, \dots, \quad (2)$$

that also satisfy the difference equation (1), are of constant sign:

$$\ell_n > 0, \quad \tilde{\ell}_n < 0, \quad n = 1, 2, \dots, \quad (3)$$

and the following limit relations hold:

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\log |\ell_n|}{n} &= \lim_{n \rightarrow \infty} \frac{\log |\tilde{\ell}_n|}{n} = \log |\mu_2| = -1.08607936 \dots, \\ \lim_{n \rightarrow \infty} \frac{\log |q_n|}{n} &= \lim_{n \rightarrow \infty} \frac{\log |p_n|}{n} = \lim_{n \rightarrow \infty} \frac{\log |\tilde{p}_n|}{n} = \log |\mu_3|, \end{aligned} \quad (4)$$

where

$$\mu_1 = -0.02001512 \dots, \quad \mu_2 = 0.33753726 \dots, \quad \mu_3 = -2368.31752213 \dots \quad (5)$$

are the roots of the characteristic polynomial $\mu^3 + 2368\mu^2 - 752\mu - 16$ of the recursion (1).

Note that the sequences $\{q_n\}$, $\{p_n\}$, and $\{\tilde{p}_n\}$ are of alternating sign:

$$(-1)^{n-1}q_n > 0, \quad (-1)^{n-1}p_n > 0, \quad (-1)^{n-1}\tilde{p}_n > 0, \quad n = 1, 2, \dots$$

Theorem 1 gives an algorithm for fast calculation of the number $\zeta(5)$. Namely, the sequence of rational numbers p_n/q_n converges to $\zeta(5)$ with speed $|\mu_2/\mu_3| < 1.42521964 \cdot 10^{-4}$ (see the table).

n	p_n/q_n	$ \zeta(5) - p_n/q_n $
0	0	$1.036927755 \dots$
1	$\frac{29}{28}$	$0.001213469 \dots$
2	$\frac{24289}{23424}$	$0.000000182 \dots$
3	$\frac{7682021239}{7408444032}$	$< 2.80 \cdot 10^{-11}$
4	$\frac{24943788950905}{24055474286592}$	$< 4.13 \cdot 10^{-15}$
5	$\frac{81875586674776013003}{78959779279372800000}$	$< 6.02 \cdot 10^{-19}$
6	$\frac{282653756112686336975107}{272587704119854963200000}$	$< 8.71 \cdot 10^{-23}$
7	$\frac{215903781003833520407770175189}{208214873150908926517286400000}$	$< 1.26 \cdot 10^{-26}$
10		$< 3.71 \cdot 10^{-38}$
20		$< 1.32 \cdot 10^{-76}$
50		$< 5.52 \cdot 10^{-192}$

Further construction of the above solutions of the difference equation (1) leads to the inclusions

$$4D_n^2 q_n \in \mathbb{Z}, \quad 4D_n^7 p_n \in \mathbb{Z}, \quad 4D_n^5 \tilde{p}_n \in \mathbb{Z}, \quad n = 1, 2, \dots$$

(see relations (14) and (15)), where D_n denotes the least common multiple of numbers $1, 2, \dots, n$, while straightforward calculations on the basis of the recursion (1) show that

$$q_n \in \mathbb{Z}, \quad 2D_n^5 p_n \in \mathbb{Z}, \quad 2D_n^3 \tilde{p}_n \in \mathbb{Z}, \quad n = 1, 2, \dots \quad (6)$$

The inclusions (6) do not allow oneself to prove the irrationality of the number $\zeta(5)$, hence we do not accent our attention on arithmetic properties of the sequences $\{q_n\}$, $\{p_n\}$, $\{\tilde{p}_n\}$ and stress only on an applied meaning of Theorem 1.

2. Auxiliary recursions. The very-well-poised hypergeometric series

$$\begin{aligned} r_n &= n!^4 \sum_{k=1}^{\infty} \left(k + \frac{n}{2}\right) \frac{\prod_{j=1}^n (k-j) \cdot \prod_{j=1}^n (k+n+j)}{\prod_{j=0}^n (k+j)^6}, \\ \tilde{r}_n &= -n!^4 \sum_{k=1}^{\infty} \left(k + \frac{n}{2}\right) \frac{\prod_{j=0}^n (k-j) \cdot \prod_{j=0}^n (k+n+j)}{\prod_{j=0}^n (k+j)^6}, \end{aligned} \quad n = 0, 1, 2, \dots, \quad (7)$$

are $\mathbb{Q}$ -linear forms in $1, \zeta(3), \zeta(5)$:

$$r_n = u_n \zeta(5) + w_n \zeta(3) - v_n, \quad \tilde{r}_n = \tilde{u}_n \zeta(5) + \tilde{w}_n \zeta(3) - \tilde{v}_n, \quad n = 0, 1, 2, \dots$$

An easy verification shows that

$$\begin{aligned} r_0 &= \zeta(5), & r_1 &= 9\zeta(5) + 33\zeta(3) - 49, & r_2 &= 469\zeta(5) + \frac{6125}{4}\zeta(3) - \frac{74463}{32}, \\ \tilde{r}_0 &= \zeta(3), & \tilde{r}_1 &= 2\zeta(5) + 12\zeta(3) - \frac{33}{2}, & \tilde{r}_2 &= 552\zeta(5) + 1764\zeta(3) - \frac{43085}{16}. \end{aligned}$$

Applying the algorithm of creative telescoping [6, Chapter 6] to the series (7) in a manner of the works [4], [5], [7], we arrive at the difference equations

$$n(n+1)^5 b_0(n-1)u_{n+1} - 2nb_1(n)u_n - b_2(n)u_{n-1} + 2(n-1)^5(2n-1)b_0(n)u_{n-2} = 0, \quad (8)$$

where

$$b_0(n) = -a_0(-n), \quad b_1(n) = a_2(-n), \quad b_2(n) = -a_1(-n),$$

and

$$n^3(n+1)^3 \tilde{b}_0(n-1)u_{n+1} - 2n\tilde{b}_1(n)u_n - \tilde{b}_2(n)u_{n-1} + 2n(n-1)^4(2n-3)\tilde{b}_0(n)\tilde{u}_{n-2} = 0, \quad (9)$$

where

$$\begin{aligned} \tilde{b}_0(n) &= 41218n^7 + 35648n^6 - 932n^5 - 13190n^4 - 5128n^3 + 811n^2 + 957n + 174, \\ \tilde{b}_1(n) &= 3874492n^{12} - 14084302n^{11} + 12425954n^{10} + 8641603n^9 - 15230839n^8 - 1369195n^7 \\ &\quad + 8618417n^6 - 623249n^5 - 2785973n^4 + 308165n^3 + 495325n^2 - 40670n - 37632, \\ \tilde{b}_2(n) &= 2(48802112n^{13} - 201803328n^{12} + 267014032n^{11} - 69927236n^{10} \\ &\quad - 95912858n^9 + 37524471n^8 + 30257812n^7 - 9523224n^6 - 8524312n^5 \\ &\quad + 2138687n^4 + 1507490n^3 - 398634n^2 - 111012n + 33408). \end{aligned}$$

Note that each of the recursions (8) and (9) has the same characteristic polynomial $\lambda^3 - 188\lambda^2 - 2368\lambda + 4$; its roots $\lambda_1, \lambda_2, \lambda_3$ ordered in increasing order of their moduli are related to the roots (5) as follows:

$$\mu_1 = \lambda_1 \lambda_2, \quad \mu_2 = \lambda_1 \lambda_3, \quad \mu_3 = \lambda_2 \lambda_3.$$

Theorem 2. *The sequences of the linear forms $\{r_n\}$ and their coefficients $\{u_n\}$, $\{w_n\}$, $\{v_n\}$ satisfy the difference equation (8). In addition, the inequalities*

$$r_n > 0, \quad u_n > 0, \quad w_n > 0, \quad v_n > 0, \quad n = 1, 2, \dots, \quad (10)$$

and the limit relations

$$\lim_{n \rightarrow \infty} \frac{\log |r_n|}{n} = \log \lambda_1, \quad \lim_{n \rightarrow \infty} \frac{\log |u_n|}{n} = \lim_{n \rightarrow \infty} \frac{\log |w_n|}{n} = \lim_{n \rightarrow \infty} \frac{\log |v_n|}{n} = \log \lambda_3 \quad (11)$$

hold.

Theorem 3. *The sequences of the linear forms $\{\tilde{r}_n\}$ and their coefficients $\{\tilde{u}_n\}$, $\{\tilde{w}_n\}$, $\{\tilde{v}_n\}$ satisfy the difference equation (9). In addition, the inequalities*

$$\tilde{r}_n < 0, \quad \tilde{u}_n > 0, \quad \tilde{w}_n > 0, \quad \tilde{v}_n > 0, \quad n = 1, 2, \dots, \quad (12)$$

and the limit relations

$$\lim_{n \rightarrow \infty} \frac{\log |\tilde{r}_n|}{n} = \log \lambda_1, \quad \lim_{n \rightarrow \infty} \frac{\log |\tilde{u}_n|}{n} = \lim_{n \rightarrow \infty} \frac{\log |\tilde{w}_n|}{n} = \lim_{n \rightarrow \infty} \frac{\log |\tilde{v}_n|}{n} = \log \lambda_3 \quad (13)$$

hold.

We mention that Vasilyev's result [8] and our theorem [4] (see also [9]) on the coincidence of very-well-poised hypergeometric series and a suitable generalization of Beukers' integral [10] lead one to the inclusions

$$\begin{aligned} 2u_n \in \mathbb{Z}, \quad 2D_n^2 w_n \in \mathbb{Z}, \quad 2D_n^5 v_n \in \mathbb{Z}, \\ 2\tilde{u}_n \in \mathbb{Z}, \quad 2D_n^2 \tilde{w}_n \in \mathbb{Z}, \quad 2D_n^5 \tilde{v}_n \in \mathbb{Z}, \end{aligned} \quad n = 1, 2, \dots \quad (14)$$

Finally, the required sequences (2) are determined by the formulae

$$\begin{aligned} \ell_n &= \tilde{w}_n r_n - w_n \tilde{r}_n = (u_n \tilde{w}_n - \tilde{u}_n w_n) \zeta(5) - (\tilde{w}_n v_n - w_n \tilde{v}_n), \\ \tilde{\ell}_n &= u_n \tilde{r}_n - \tilde{u}_n r_n = (u_n \tilde{w}_n - \tilde{u}_n w_n) \zeta(3) - (u_n \tilde{v}_n - \tilde{u}_n v_n), \end{aligned} \quad n = 0, 1, 2, \dots,$$

hence

$$q_n = u_n \tilde{w}_n - \tilde{u}_n w_n, \quad p_n = \tilde{w}_n v_n - w_n \tilde{v}_n, \quad \tilde{p}_n = u_n \tilde{v}_n - \tilde{u}_n v_n, \quad n = 0, 1, 2, \dots \quad (15)$$

For proving the inequalities (3) and the limit relations (4), it remains to use the estimates (10), (12), relations (11), (13), and Poincaré's theorem.

3. Conclusion remarks. Recently, another algorithm for fast calculation of $\zeta(5)$ and $\zeta(3)$ (also not producing good enough diophantine approximations to these constants), based on the infinite matrix product

$$\prod_{n=1}^{\infty} \begin{pmatrix} -\frac{n}{2(2n+1)} & \frac{1}{2n(2n+1)} & \frac{1}{n^4} \\ 0 & -\frac{n}{2(2n+1)} & \frac{5}{4n^2} \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 & \zeta(5) \\ 0 & 0 & \zeta(3) \\ 0 & 0 & 1 \end{pmatrix},$$

has been suggested by R. W. Gosper [11]. However the letters [11] do not contain any description of analytic and arithmetic properties of the rational approximations so constructed. Mention also an algorithm due to E. A. Karatsuba [12] for calculation of the Riemann zeta function at positive integers.

The above scheme of Section 2 also allows one to construct a third-order recursion for simultaneous rational approximations to $\zeta(2)$ and $\zeta(3)$. Namely, consider the difference equation

$$(n+1)^4 \tilde{a}_0(n) q_{n+1} - \tilde{a}_1(n) q_n + 4(2n-1) \tilde{a}_2(n) q_{n-1} - 4(n-1)^2 (2n-1)(2n-3) \tilde{a}_0(n+1) q_{n-2} = 0, \quad (16)$$

where

$$\begin{aligned} \tilde{a}_0(n) &= 946n^2 - 731n + 153, \\ \tilde{a}_1(n) &= 2(104060n^6 + 127710n^5 + 12788n^4 - 34525n^3 - 8482n^2 + 3298n + 1071), \\ \tilde{a}_2(n) &= 3784n^5 - 1032n^4 - 1925n^3 + 853n^2 + 328n - 184, \end{aligned}$$

and define its three linearly independent solutions $\{q'_n\}$, $\{p'_n\}$, and $\{\tilde{p}'_n\}$ by the initial data

$$q'_0 = 1, \quad q'_1 = 14, \quad q'_2 = 978, \quad p'_0 = 0, \quad p'_1 = 17, \quad p'_2 = \frac{9405}{8}, \quad \tilde{p}'_0 = 0, \quad \tilde{p}'_1 = 23, \quad \tilde{p}'_2 = \frac{6435}{4}.$$

Theorem 4. *The sequences $\{q'_n\}$, $\{p'_n\}$, and $\{\tilde{p}'_n\}$ (of positive sign) as well as the sequences*

$$\ell'_n = q'_n \zeta(3) - p'_n, \quad \tilde{\ell}'_n = q'_n \zeta(2) - \tilde{p}'_n, \quad n = 0, 1, 2, \dots,$$

satisfy the limit relations

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\log |\ell'_n|}{n} &= \lim_{n \rightarrow \infty} \frac{\log |\tilde{\ell}'_n|}{n} = \log |\mu_2| = -1.31018925\dots, \\ \lim_{n \rightarrow \infty} \frac{\log |q'_n|}{n} &= \lim_{n \rightarrow \infty} \frac{\log |p'_n|}{n} = \lim_{n \rightarrow \infty} \frac{\log |\tilde{p}'_n|}{n} = \log |\mu_3|, \end{aligned}$$

where

$$\mu_{1,2} = 0.07260980\dots \pm i0.25981363\dots, \quad \mu_3 = 219.85478039\dots$$

are the roots of the characteristic polynomial $\mu^3 - 220\mu^2 + 32\mu - 16$ of the recursion (16).

This time, auxiliary recursions are satisfied by the hypergeometric series

$$r'_n = -n!^2 \sum_{k=1}^{\infty} \frac{\prod_{j=1}^n (k-j)}{\prod_{j=0}^n (k+j)^3}, \quad \tilde{r}'_n = n!^2 \sum_{k=1}^{\infty} \frac{\prod_{j=0}^n (k-j)}{\prod_{j=0}^n (k+j)^3}, \quad n = 0, 1, 2, \dots,$$

that are $\mathbb{Q}$ -linear forms in 1, $\zeta(2)$, and $\zeta(3)$.

Similarly to the case of the difference equation (1), the explicit calculations lead to the inclusions

$$q'_n \in \mathbb{Z}, \quad D_n^3 p'_n \in \mathbb{Z}, \quad D_n^2 \tilde{p}'_n \in \mathbb{Z}, \quad n = 1, 2, \dots,$$

that are quite better than one can expect.

REFERENCES

1. R. Apéry, *Irrationalité de $\zeta(2)$ et $\zeta(3)$* , Astérisque **61** (1979), 11–13.
2. A. van der Poorten, *A proof that Euler missed... Apéry's proof of the irrationality of $\zeta(3)$* , An informal report, Math. Intelligencer **1** (1978/79), no. 4, 195–203.
3. V. N. Sorokin, *One algorithm for fast calculation of $\zeta(4)$* , Preprint (April 2002), Russian Academy of Sciences, M. V. Keldysh Institute for Applied Mathematics, Moscow, 2002.
4. W. Zudilin, *Well-poised hypergeometric service for diophantine problems of zeta values*, Actes des 12èmes rencontres arithmétiques de Caen (June 29–30, 2001), J. Théorie Nombres Bordeaux (2003) (to appear).
5. W. Zudilin, *Apéry-like difference equation for Catalan's constant*, E-print math.NT/0201024 (January 2002).
6. M. Petkovšek, H. S. Wilf, and D. Zeilberger, *A = B*, A. K. Peters, Ltd., Wellesley (M.A.), 1997.
7. W. Zudilin, *An elementary proof of Apéry's theorem*, E-print math.NT/0202159 (February 2002).
8. D. V. Vasilyev, *On small linear forms for the values of the Riemann zeta-function at odd points*, Preprint no. 1 (558), Nat. Acad. Sci. Belarus, Institute Math., Minsk, 2001.
9. W. Zudilin, *Very-well-poised hypergeometric series and multiple integrals*, Uspekhi Mat. Nauk [Russian Math. Surveys] **57** (2002), no. 4; E-print math.CA/0206177 (March 2002).
10. F. Beukers, *A note on the irrationality of $\zeta(2)$ and $\zeta(3)$* , Bull. London Math. Soc. **11** (1979), no. 3, 268–272.
11. R. W. Gosper, *Some infinite products involving $\zeta(2m+1)$* , Letters of 24.10.2000 and 25.10.2000, Favorite Mathematical Constants (S. Finch, ed.), <http://pauillac.inria.fr/algo/bsolve/constant/apery/infprd.html>, 2000.
12. E. A. Karatsuba, *Fast computation of the Riemann zeta function $\zeta(s)$ for integer values of s* , Problemy Peredachi Informatsii [Problems Inform. Transmission] **31** (1995), no. 4, 69–80.

MOSCOW LOMONOSOV STATE UNIVERSITY

DEPARTMENT OF MECHANICS AND MATHEMATICS

VOROBIOVY GORY, GSP-2, 119992 MOSCOW, RUSSIA

URL: <http://wain.mi.ras.ru/index.html>

E-mail address: wadim@ips.ras.ru